VI Autonomous Agents & Self-Organization

2012/4/4

Autonomous Agent

- "a unit that interacts with its environment (which probably consists of other agents)
- but acts independently from all other agents in that it does not take commands from some seen or unseen leader,
- nor does an agent have some idea of a global plan that it should be following."
 Flake (p. 261)





A. Schools, Flocks, & Herds

"and the thousands of fishes moved as a huge beast, piercing the water. They appeared united, inexorably bound to a common fate. How comes this unity?"

- anon., 17th cent.

2012/4/4

images from EVALife site

3

Coordinated Collective Movement

- Groups of animals can behave almost like a single organism
- Can execute swift maneuvers
 - for predation or to avoid predation
- Individuals rarely collide, even in frenzy of attack or escape
- Shape is characteristic of species, but flexible


Adaptive Significance


- Prey avoiding predation
- More efficient predation by predators
- Other efficiencies


2012/4/4


Avoiding Predation


- More compact aggregation
 - predator risks injury by attacking
- Confusing predator by:
 - united erratic maneuvers (e.g. zigzagging)
 - separation into subgroups (e.g., flash expansion & fountain effect)














Better Predation

- Coordinated movements to trap prey
 - e.g. parabolic formation of tuna
- More efficient predation
 - e.g., killer whales encircle dolphins
 - take turns eating

2012/4/4

Other Efficiencies

- Fish schooling may increase hydrodynamic efficiency
 - endurance may be increased up to 6x
 - school acts like "group-level vehicle"
- V-formation increases efficiency of geese
 - range 70% greater than that of individual
- Lobsters line up single file by touch
 - move 40% faster than when isolated
 - decreased hydrodynamic drag

Characteristic Arrangement of School

- Shape is characteristic of species
- Fish have preferred distance, elevation & bearing relative to neighbors
- Fish avoid coming within a certain minimum distance
 - closer in larger schools
 - closer in faster moving schools

2012/4/4

Alternatives to Self-Organization

- "Templates"
 - no evidence that water currents, light, chemicals guide collective movement
- "Leaders"
 - no evidence for leaders
 - those in front may drop behind
 - those on flank may find selves in front
 - each adjusts to several neighbors
- "Blueprint" or "Recipe"
 - implausible for coordination of large schools
 - e.g., millions of herring, hundreds of millions of cod

Self-Organization Hypothesis

- Simple attraction & repulsion rules generate schooling behavior
 - positive feedback: brings individuals together
 - negative feedback: but not too close
- Rules rely on local information
 - i.e. positions & headings of a few nearby fish
 - no global plan or centralized leader

2012/4/4

Mechanisms of Individual Coordination

- Vision
 - governs attraction
 - & alignment
- Lateral line
 - sensitive to water movement
 - provides information on speed & direction of neighbors
 - governs repulsion
 - & speed matching
- How is this information integrated into a behavioral plan?
 - most sensitive to nearest neighbors

Basic Assumptions of Huth & Wissel (1992) Model

- All fish follow same rules
- Each uses some sort of weighted average of positions & orientations of nearest neighbors
- Fish respond to neighbors probabilistically
 - imperfect information gathering
 - imperfect execution of actions
- No external influences affect fish
 - e.g. no water currents, obstacles, ...

2012/4/4

Ranges of Behavior Patterns attract orient search search 2012/4/4 Fig. adapted from Camazine & al., Self-Org. Biol. Sys. 20

Model Behavior of Individual

1. Determine a target direction from each of three nearest neighbors:

if in repel range, then 180° + direction to neighbor else if in orient range, then heading of neighbor else if in attract range, then accelerate if ahead, decelerate if behind; return direction to neighbor

else return our own current heading

- 2. Determine overall target direc. as average of 3 neighbors inversely weighted by their distances
- 3. Turn a fraction in this direction (determined by *flexibility*) + some randomness

2012/4/4

Demonstration of NetLogo Simulation of Flocking/ Schooling based on Huth & Wissel Model

Run Flock.nlogo

Limitations of Model

- Model addresses only motion in absence of external influences
- Ignores obstacle avoidance
- Ignores avoidance behaviors such as:
 - flash expansion
 - fountain effect
- Recent work (since 1997) has addressed some of these issues

2012/4/4 23

"Boids"

A model of flocks, herds, and similar cases of coordinated animal motion by Craig Reynolds (1986)

NetLogo Simulation

- Flockmates are those within "vision"
- If nearest flockmate < minimum separation, turn away
- Else:
 - align with average heading of flockmates
 - cohere by turning toward average flockmate direction
- All turns limited specified maxima
- Note fluid behavior from deterministic rules

2/4/4

Demonstration of boids

Run Flocking.nlogo

Demonstration of boids (with 3D perspective)

Run Flocking (Perspective Demo).nlogo

2012/4/4

Demonstration of 3D boids

Run Flocking 3D.nlogo

Obstacle Avoidance

- Boid flock avoiding cylindrical obstacles (Reynolds 1986)
- This model incorporates:
 - predictive obstacle avoidance
 - goal seeking (scripted path)

COURSE: 07
COURSE ORGANIZER: DEMETRI TERZODOULOS
"BOIDS DEMOS"
CRAIG REVNOLDS
SILICON STUDIOS, MS 3L-080
2011 NORTH SHORELINE BLVD.
MOUNTAIN VIEW, (A 94039-7311

2012/4/4

29

Jon Klein's Flocking Algorithm

- Sight limited by "vision"
- Balances 6 "urges":
 - be near center of flock
 - have same velocity as flockmates
 - keep spacing correct
 - avoid collisions with obstacles
 - be near center of world
 - wander throughout world
- Strength of urge affects acceleration

Demonstration of Klein's Flocking Algorithm

Run Flocking 3D Alternate.nlogo

2012/4/4

Use in Computer Animation



- Extract from Stanley and Stella in "Breaking the Ice" (1987)
- store.yahoo.com/ odyssey3d/ comanclascli2.html


Particle Swarm Optimization


(Kennedy & Eberhart, 1995)


2012/4/4


Motivation


- Originally a model of social information sharing
- Abstract vs. concrete spaces
 - cannot occupy same locations in concrete space
 - can in abstract space (two individuals can have same idea)
- Global optimum (& perhaps many suboptima)
- Combines:
 - private knowledge (best solution each has found)
 - public knowledge (best solution entire group has found)























Variables

 $\mathbf{x}_k = \text{current position of particle } k$

 \mathbf{v}_k = current velocity of particle k

 \mathbf{p}_k = best position found by particle k

 $Q(\mathbf{x})$ = quality of position \mathbf{x}

g = index of best position found so far

i.e., $g = \operatorname{argmax}_k Q(\mathbf{p}_k)$

 ϕ_1 , ϕ_2 = random variables uniformly distributed over [0, 2]

w = inertia < 1

2012/4/4

44

Velocity & Position Updating

 $\mathbf{v}_{k}' = w \ \mathbf{v}_{k} + \phi_{1} \ (\mathbf{p}_{k} - \mathbf{x}_{k}) + \phi_{2} \ (\mathbf{p}_{g} - \mathbf{x}_{k})$ $w \ \mathbf{v}_{k}$ maintains direction (*inertial* part) $\phi_{1} \ (\mathbf{p}_{k} - \mathbf{x}_{k})$ turns toward private best (*cognition* part) $\phi_{2} \ (\mathbf{p}_{g} - \mathbf{x}_{k})$ turns towards public best (*social* part)

 $\mathbf{x}_{k}' = \mathbf{x}_{k} + \mathbf{v}_{k}'$

- Allowing $\phi_1, \phi_2 > 1$ permits overshooting and better exploration (*important!*)
- Good balance of exploration & exploitation
- Limiting $\|\mathbf{v}_k\| < \|\mathbf{v}_{\text{max}}\|$ controls resolution of search

2012/4/4

Netlogo Demonstration of Particle Swarm Optimization

Run PSO.nlogo

or Particle Swarm Optimization.nlogo

Yuhui Shi's Demonstration of Particle Swarm Optimization

Run www.engr.iupui.edu/~shi/PSO/ AppletGUI.html

2012/4/4


Improvements

- Alternative velocity update equation:
 - $\mathbf{v}_{k}' = \mathbf{\chi} \left[w \mathbf{v}_{k} + \mathbf{\phi}_{1} \left(\mathbf{p}_{k} \mathbf{x}_{k} \right) + \mathbf{\phi}_{2} \left(\mathbf{p}_{g} \mathbf{x}_{k} \right) \right]$

 χ = constriction coefficient (controls magnitude of \mathbf{v}_k)

- Alternative neighbor relations:
 - star: fully connected (each responds to best of all others; fast information flow)
 - circle: connected to K immediate neighbors (slows information flow)
 - wheel: connected to one axis particle (moderate information flow)

Spatial Extension



- Spatial extension avoids premature convergence
- Preserves diversity in population
- More like flocking/schooling models

2012/4/4 Fig. from EVALife site

Netlogo Demonstration of Particle Swarm Optimization with Collision Avoidance

Run PSO.nlogo

Some Applications of PSO

- integer programming
- minimax problems
 - in optimal control
 - engineering design
 - discrete optimization
 - Chebyshev approximation
 - game theory
- multiobjective optimization
- hydrologic problems
- musical improvisation!

2012/4/4

Millonas' Five Basic Principles of Swarm Intelligence

- 1. Proximity principle:
 pop. should perform simple space & time computations
- 2. Quality principle: pop. should respond to quality factors in environment
- 3. Principle of diverse response: pop. should not commit to overly narrow channels
- 4. Principle of stability:

 pop. should not change behavior every time env. changes
- 5. Principle of adaptability: pop. should change behavior when it's worth comp. price

2012/4/4 (Millonas 1994) 52

Kennedy & Eberhart on PSO

"This algorithm belongs ideologically to that philosophical school

that allows wisdom to emerge rather than trying to impose it,

that emulates nature rather than trying to control it, and that seeks to make things simpler rather than more complex.

Once again nature has provided us with a technique for processing information that is at once elegant and versatile."

2012/4/4 53

Additional Bibliography

- 1. Camazine, S., Deneubourg, J.-L., Franks, N. R., Sneyd, J., Theraulaz, G., & Bonabeau, E. *Self-Organization in Biological Systems*. Princeton, 2001, chs. 11, 13, 18, 19.
- 2. Bonabeau, E., Dorigo, M., & Theraulaz, G. Swarm Intelligence: From Natural to Artificial Systems. Oxford, 1999, chs. 2, 6.
- 3. Solé, R., & Goodwin, B. Signs of Life: How Complexity Pervades Biology. Basic Books, 2000, ch. 6.
- 4. Resnick, M. *Turtles, Termites, and Traffic Jams: Explorations in Massively Parallel Microworlds*. MIT Press, 1994, pp. 59-68, 75-81.
- 5. Kennedy, J., & Eberhart, R. "Particle Swarm Optimization," *Proc. IEEE Int'l. Conf. Neural Networks* (Perth, Australia), 1995. http://www.engr.iupui.edu/~shi/pso.html.

2012/4/4



54