

Project 2

Creation of Spatial Structure by Activator/Inhibitor Cellular Automaton

Due February 18, 2015

Background

Activator/Inhibitor Cellular Automaton

In this project you will investigate and measure the creation of spatial structure by an activator/inhibitor CA (AICA) such as we discussed in class.¹ Structure will be quantified in terms of *spatial correlation* and *mutual information* (discussed below), which you will measure after the CA has converged to a stable state.

Recall that the state transition rule of an activator/inhibitor network is given by a formula such as this:

$$s_i(t+1) = \text{sign} \left[h + J_1 \sum_{r_{ij} < R_1} s_j(t) + J_2 \sum_{R_1 \leq r_{ij} < R_2} s_j(t) \right]. \quad (1)$$

Since this is a 2D CA, the cell indices are two-dimensional vectors, $\mathbf{i} = (i_1, i_2)$, $\mathbf{j} = (j_1, j_2)$. As usual, we will also assume that the space is a torus, that is, the indices wrap around on the top and bottom and on the left and right edges. For the purposes of this project you may assume that $J_1 \geq 0$ and $J_2 \leq 0$ (which are the usual cases). Also, you should assume that the R_1 neighborhood includes the cell (\mathbf{i}) at its center.

The distance r_{ij} between cells can be measured in many different ways, for example by Euclidean distance (also known as the L_2 metric). For the purpose of this project it will be more convenient to measure distance by the L_1 metric, which is defined:

$$r_{ij} = |i_1 - j_1| + |i_2 - j_2|. \quad (2)$$

This gives neighborhoods that are diamond-shaped, rather than circular, but that doesn't really matter.

¹Additional information and tips are available on Dr. Van Hornweder's help page:
<<http://web.eecs.utk.edu/~mclennan/Classes/420/projects/project2/aica.html>>.

Spatial Correlation

We are interested in the extent to which the states of cells at various distances are correlated to each other. Therefore, we define the *absolute spatial correlation at distance l* as

$$\rho_l = |\langle s_i s_j \rangle - \langle s_i \rangle \langle s_j \rangle| \text{ for all } \mathbf{i}, \mathbf{j} \text{ such that } r_{\mathbf{ij}} = l.^2$$

The angle brackets mean “average value of.” To understand this formula, suppose that the states $+1$ and -1 are equally frequent in the space. Then the individual state averages are 0: $\langle s_i \rangle = \langle s_j \rangle = 0$. Therefore $\langle s_i s_j \rangle$ is the average value of the product $s_i s_j$ for cells that are a distance l apart. If these cells tend to be $+1$ at the same time, or -1 at the same time, this average will be greater than zero (positive correlation). If they tend to have opposite signs, then it will be less than zero (negative correlation). If they tend to have the same sign as often as the opposite sign, then the average will be near zero (no correlation). By subtracting $\langle s_i \rangle \langle s_j \rangle$ we compensate for an overall bias toward positive or negative states (such as we get when $h \neq 0$). We take the absolute value, because we are not interested in whether the spatial correlation is positive or negative, only its magnitude. Note that $\rho_0 = 1 - \langle s_i \rangle^2$.

Next let's consider more explicitly how to compute ρ_l . Suppose there are N^2 cells in the space, and let C_l ($1 \leq l < N/2$) be the circumference (in number of cells) of a neighborhood of radius l . For the L_1 metric, $C_l = 4l$. Thus there are C_l cells at a distance l from a given cell. Then we can see (make sure you really do see it!) that:

$$\rho_l = \left| \frac{2}{N^2 C_l} \sum_{\substack{\langle \mathbf{ij} \rangle \\ r_{\mathbf{ij}} = l}} s_i s_j - \left(\frac{1}{N^2} \sum_{\mathbf{i}} s_i \right)^2 \right|.$$

The notation $\langle \mathbf{ij} \rangle$ under the summation means “all pairs of distinct cells \mathbf{i} and \mathbf{j} ” (taking each pair just once); therefore there are $N^2 C_l / 2$ of these pairs. (There is a diamond of C_l cells around each of the N^2 cells.) Thus we are averaging over all pairs at a distance of l . For purposes of computation, this can be made more explicit:

$$\rho_l = \left| \left(\frac{1}{4lN^2} \sum_{i_1, i_2} \left(s_{i_1, i_2} \sum_{|j_1| + |j_2| = l} s_{i_1 + j_1, i_2 + j_2} \right) \right) - \left(\frac{1}{N^2} \sum_{i_1, i_2} s_{i_1, i_2} \right)^2 \right|.$$

Notice that the second summation is over positive and negative j_1, j_2 in the

² Note: A true correlation coefficient is normalized to $[-1, 1]$ by dividing by standard deviations of the variables: $\rho_l = \left| \left(\langle s_i s_j \rangle - \langle s_i \rangle \langle s_j \rangle \right) / \sigma_{s_i} \sigma_{s_j} \right|$ for all \mathbf{i}, \mathbf{j} such that $r_{\mathbf{ij}} = l$. You can compute it this way if you want, but make sure to tell us.

range $-l \leq j_1, j_2 \leq l$. The coefficient on the first summation is:

$$\frac{1}{2} \frac{2}{N^2} \frac{1}{C_l} = \frac{1}{4lN^2}.$$

The $\frac{1}{2}$ factor is to avoid double counting the pairs $\langle \mathbf{ij} \rangle$. Make sure that you understand the formula for ρ_l . Remember that cell indices wrap around both vertically and horizontally.

Characteristic Correlation Length

(This section is relevant to **required measurements for COSC 427 and 527 students**. COSC 420 students can skip it, unless they want to do these measurements for extra credit.)

Often spatial correlation decreases approximately exponentially with distance, $\rho_l \propto e^{-l/\lambda}$, where λ , the *characteristic correlation length*, measures how quickly spatial correlation decreases with distance. By assuming that spatial correlation is exponentially decreasing, we can estimate λ . Since ρ_0 is special, we compute it relative to ρ_1 . Let α be an arbitrary constant of proportionality $\rho_l = \alpha e^{-(l-1)/\lambda}$. Then,

$$\begin{aligned} \rho_1 &= \alpha e^0 = \alpha, \\ \rho_{\lambda+1} &= \alpha e^{-\lambda/\lambda} = \alpha e^{-1} = \frac{\rho_1}{e}. \end{aligned}$$

Therefore we can estimate $\lambda \approx l - 1$ such that $\rho_l = \rho_1/e$. That is, the characteristic correlation length is that length at which the spatial correlation has decreased to $1/e$ of its hypothetical maximum value ρ_1 . (*Characteristic correlation time* can be defined similarly.)

Mutual Information Between Distant Cells

As discussed in class, one way to measure the correlation between cell states is by *average mutual information*. The average mutual information between cells at a distance l is related to the joint entropy between cells at this distance.

Therefore, first define the average entropy $H(S)$ of the cellular space S :

$$H(S) = - \sum_{s \in \{-1, +1\}} \Pr\{s\} \lg \Pr\{s\}.$$

Remember that we take $0 \lg 0 = 0$. For state counting convenience, let $\beta(s) = (1+s)/2$ so that $\beta(+1) = 1$ and $\beta(-1) = 0$ (β converts a *bipolar* number $\in \{-1, +1\}$ into a *binary* number $\in \{0, 1\}$). Then the probabilities of the states can be computed:

$$\begin{aligned} \Pr\{+1\} &= \frac{1}{N^2} \sum_i \beta(s_i), \\ \Pr\{-1\} &= 1 - \Pr\{+1\}. \end{aligned}$$

The probability that two cells at distance l have state $+1$ is:

$$P_l(+1, +1) = \frac{2}{N^2 C_l \langle ij \rangle} \sum_{r_{ij}=l} \beta(s_i) \beta(s_j) .$$

Similarly,

$$P_l(-1, -1) = \frac{2}{N^2 C_l \langle ij \rangle} \sum_{r_{ij}=l} \beta(-s_i) \beta(-s_j) .$$

These summations can be computed in the same way as ρ_l . Finally,

$$P_l(+1, -1) = P_l(-1, +1) = 1 - P_l(+1, +1) - P_l(-1, -1) .$$

(Notice that there are only three distinct possibilities for cells at distance l : both $+1$, both -1 , or opposite signs.) The joint entropy between cells at a distance l , H_l , is then defined in terms of the probabilities in the usual way:

$$H_l = - \left(P_l(+1, +1) \lg P_l(+1, +1) + P_l(-1, -1) \lg P_l(-1, -1) + P_l(+1, -1) \lg P_l(+1, -1) \right) .$$

The *average mutual information* between two sources A and B is defined $I(A, B) = H(A) + H(B) - H(A, B)$. Therefore, the average mutual information between cells at distance l is defined:

$$I_l = 2H(S) - H_l .$$

Experiments

In this project you will be investigating how spatial structure (as measured by spatial correlation and mutual information) is affected by the parameters (J_1 , J_2 , R_1 , R_2 , h) of an AICA. (Graduate students should also compute characteristic correlation length λ ; undergraduates can do this for extra credit.) Make sure to check Dr. Van Hornweder's help page for information and suggestions on doing your project:

<<http://web.eecs.utk.edu/~mclennan/Classes/420/projects/project2/aica.html>>.

The goal is to get the measures of structure for a variety of different combinations of parameters for each of the following three experiments. The following table shows the minimum set of parameter combinations that you should use. You can try additional combinations to earn some extra credit.

Your CA space should be $N=30$ in each dimension. Use asynchronous updating (i.e., update the cells one at a time rather than all at once), and update the states randomly, not in a systematic order, or you will get artifacts.

For each set of parameters you will make several runs, with different random initial states, so that you can average your results. Do a few trial runs to see how much the measures vary from run to run. If they don't vary much, then you don't need to do many runs for each set of parameters (maybe one run will be enough). If they do vary, then you will have to make several runs (less than 5) and average them together, and you won't have to investigate so many parameter values.

For each run you should allow the CA to stabilize, and then compute $H(S)$ estimated λ (this is optional for COSC 420), and also plot I_l and ρ_l against the l values. For each set of parameter values, compute the average of all of these and print them out (generate plots for I_l and ρ_l). You don't need to hand in a graph for every set of parameters, just those that are relevant to your hypotheses.

Following are the descriptions of specific experiments that you should try. In each case, discuss and explain your results.

Descriptions of Experiments:

1. **(Extra Credit)** Set $J_2=0$ to disable the inhibition system and set $J_1=1$. Quantify the spatial structure for at least the specified range of R_1 and h values.
2. **(Extra Credit)** Set $J_1=0$ to disable the activation system and set $J_2=-0.1$. Quantify spatial structure as in Experiment (1) for at least the specified parameter values.
3. Set $J_1=1$ and $J_2=-0.1$. Investigate, as in the preceding experiments, the spatial structure for at least the specified values of R_1 , R_2 , and h .

Based on all the preceding experiments, draw conclusions about the dependence of spatial correlation and mutual information on the parameters of the AICA. See if you can draw some quantitative conclusions.

| Experiment 1 (extra credit) | | | Experiment 2 (extra credit) | | | Experiment 3 (required) | | |
|--------------------------------|-------|-----|--------------------------------|-------|-----|----------------------------|-------|-----|
| R_1 | R_2 | h | R_1 | R_2 | h | R_1 | R_2 | h |
| 1 | 15 | -1 | 1 | 2 | 0 | 1 | 2 | 0 |
| 3 | 15 | -1 | 1 | 4 | -2 | 1 | 5 | -4 |
| 6 | 15 | -2 | 1 | 4 | -1 | 1 | 5 | -2 |
| | | | 1 | 4 | 0 | 1 | 5 | 0 |
| | | | 1 | 6 | -5 | 1 | 9 | -6 |
| | | | 1 | 6 | -3 | 1 | 9 | -3 |
| | | | 1 | 6 | 0 | 1 | 9 | 0 |
| | | | 1 | 9 | 0 | 1 | 14 | 0 |
| | | | 1 | 13 | 0 | 3 | 5 | -1 |
| | | | 4 | 5 | 0 | 3 | 5 | 0 |
| | | | 4 | 7 | -5 | 3 | 9 | -6 |
| | | | 4 | 7 | -3 | 3 | 9 | -3 |
| | | | 4 | 7 | 0 | 3 | 9 | 0 |
| | | | 4 | 12 | 0 | 3 | 14 | 0 |
| | | | 9 | 12 | -6 | 7 | 9 | 0 |
| | | | 9 | 12 | -3 | 7 | 14 | 0 |
| | | | 9 | 12 | 0 | 12 | 14 | 0 |