

V. Evolutionary Computing

A. Genetic Algorithms

11/2/09

1

Genetic Algorithms

- Developed by John Holland in '60s
- Did not become popular until late '80s
- A simplified model of genetics and evolution by natural selection
- Most widely applied to optimization problems (maximize "fitness")

11/2/09

2

Assumptions

- Existence of fitness function to quantify merit of potential solutions
 - this "fitness" is what the GA will maximize
- A mapping from bit-strings to potential solutions
 - best if each possible string generates a legal potential solution
 - choice of mapping is important
 - can use strings over other finite alphabets

11/2/09

3

Outline of Simplified GA

1. Random initial population $P(0)$
2. Repeat for $t = 0, \dots, t_{\max}$ or until converges:
 - a) create empty population $P(t + 1)$
 - b) repeat until $P(t + 1)$ is full:
 - 1) select two individuals from $P(t)$ based on fitness
 - 2) optionally mate & replace with offspring
 - 3) optionally mutate offspring
 - 4) add two individuals to $P(t + 1)$

11/2/09

4

Fitness-Biased Selection


- Want the more "fit" to be more likely to reproduce
 - always selecting the best
⇒ premature convergence
 - probabilistic selection ⇒ better exploration
- Roulette-wheel selection: probability \propto relative fitness:

$$\Pr\{i \text{ mates}\} = \frac{f_i}{\sum_{j=1}^n f_j}$$

11/2/09

5

Crossover: Biological Inspiration






- Occurs during meiosis, when haploid gametes are formed
- Randomly mixes genes from two parents
- Creates genetic variation in gametes

11/2/09

(fig. from B&N Thes., Biol.)

6



Mutation: Biological Inspiration

- **Chromosome mutation** ⇒
- **Gene mutation:** alteration of the DNA in a gene
 - inspiration for mutation in GAs
- In typical GA each bit has a low probability of changing
- Some GAs models rearrange bits

(fig. from *B&N Thes. Biol.*)

11/2/09 10

The Red Queen Hypothesis

- **Observation:** a species probability of extinction is independent of time it has existed
- **Hypothesis:** species continually adapt to each other
- Extinction occurs with insufficient variability for further adaptation

“Now, *here*, you see, it takes all the running *you* can do, to keep in the same place.”
 — *Through the Looking-Glass and What Alice Found There*

11/2/09 11

Demonstration of GA: Finding Maximum of Fitness Landscape

[Run Genetic Algorithms — An Intuitive Introduction](#)
 by [Pascal Glauser](#)
www.glauserweb.ch/gentore.htm

11/2/09 12

Demonstration of GA: Evolving to Generate a Pre-specified Shape (Phenotype)

[Run Genetic Algorithm Viewer
<www.rennard.org/alife/english/gavgb.html>](http://www.rennard.org/alife/english/gavgb.html)

11/2/09

13

Demonstration of GA: Eaters Seeking Food

<http://math.hws.edu/xJava/GA/>

11/2/09

14

Morphology Project by Michael "Flux" Chang

- Senior Independent Study project at UCLA
 - users.design.ucla.edu/~mflux/morphology
- Researched and programmed in 10 weeks
- Programmed in **Processing** language
 - www.processing.org

11/2/09

15

Genotype \Rightarrow Phenotype

- Cells are "grown," not specified individually
- Each gene specifies information such as:
 - angle
 - distance
 - type of cell
 - how many times to replicate
 - following gene
- Cells connected by "springs"
- Run **phenome**:
 - users.design.ucla.edu/~mflux/morphology/gallery/sketches/phenome

11/2/09

16

Complete Creature

- Neural nets for control (**blue**)
 - integrate-and-fire neurons
- Muscles (**red**)
 - decrease "spring length" when fire
- Sensors (**green**)
 - fire when exposed to "light"
- Structural elements (**grey**)
 - anchor other cells together
- Creature embedded in a fluid
- Run users.design.ucla.edu/~mflux/morphology/gallery/sketches/creature

11/2/09

17

Effects of Mutation

- Neural nets for control (**blue**)
- Muscles (**red**)
- Sensors (**green**)
- Structural elements (**grey**)
- Creature embedded in a fluid
- Run
 - users.design.ucla.edu/~mflux/morphology/gallery/sketches/creaturepack

11/2/09

18


Evolution

- Population: 150–200
- Nonviable & non-responsive creatures eliminated
- Fitness based on speed or light-following
- 30% of new pop. are mutated copies of best
- 70% are random
- No crossover



11/2/09 19

Gallery of Evolved Creatures



- Selected for speed of movement
- Run

users.design.ucla.edu/~mflux/morphology/gallery/sketches/creaturegallery

11/2/09 20


Why Does the GA Work?

The Schema Theorem

11/2/09 21

Schemata

A **schema** is a description of certain patterns of bits in a genetic string



11/2/09 22

The Fitness of Schemata

- The schemata are the **building blocks** of solutions
- We would like to know the average fitness of all possible strings belonging to a schema
- We cannot, but the strings in a population that belong to a schema give an estimate of the fitness of that schema
- Each string in a population is giving information about all the schemata to which it belongs (**implicit parallelism**)

11/2/09 23

Effect of Selection

Let n = size of population

Let $m(S, t)$ = number of instances of schema S at time t

String i gets picked with probability $\frac{f_i}{\sum_j f_j}$

Let $f(S)$ = avg fitness of instances of S at time t

So expected $m(S, t + 1) = m(S, t) \cdot n \cdot \frac{f(S)}{\sum_j f_j}$

Since $f_{av} = \frac{\sum_j f_j}{n}$, $m(S, t + 1) = m(S, t) \frac{f(S)}{f_{av}}$

11/2/09 24

Exponential Growth

- We have discovered:

$$m(S, t+1) = m(S, t) \cdot f(S) / f_{av}$$
- Suppose $f(S) = f_{av} (1 + c)$
- Then $m(S, t) = m(S, 0) (1 + c)^t$
- That is, **exponential growth** in above-average schemata

11/2/09 25

Effect of Crossover

$**1 \dots 0**$
 $|\leftarrow \delta \rightarrow|$

- Let λ = length of genetic strings
- Let $\delta(S)$ = defining length of schema S
- Probability {crossover destroys S }:

$$p_d \approx \delta(S) / (\lambda - 1)$$
- Let p_c = probability of crossover
- Probability schema survives:

$$p_s \geq 1 - p_c \frac{\delta(S)}{\lambda - 1}$$

11/2/09 26

Selection & Crossover Together

$$m(S, t+1) \geq m(S, t) \frac{f(S)}{f_{av}} \left[1 - p_c \frac{\delta(S)}{\lambda - 1} \right]$$

11/2/09 27

Effect of Mutation

- Let p_m = probability of mutation
- So $1 - p_m$ = probability an allele survives
- Let $o(S)$ = number of fixed positions in S
- The probability they all survive is

$$(1 - p_m)^{o(S)}$$
- If $p_m \ll 1$, $(1 - p_m)^{o(S)} \approx 1 - o(S) p_m$

11/2/09 28

Schema Theorem: “Fundamental Theorem of GAs”

$$m(S, t+1) \geq m(S, t) \frac{f(S)}{f_{av}} \left[1 - p_c \frac{\delta(S)}{\lambda - 1} - o(S) p_m \right]$$

11/2/09 29

The Bandit Problem

- Two-armed bandit:
 - random payoffs with (unknown) means m_1, m_2 and variances σ_1, σ_2
 - optimal strategy: allocate exponentially greater number of trials to apparently better lever
- k -armed bandit: similar analysis applies
- Analogous to allocation of population to schemata
- Suggests GA may allocate trials optimally

11/2/09 30

Goldberg's Analysis of Competent & Efficient GAs

11/2/09 31

Paradox of GAs

- Individually uninteresting operators:
 - selection, recombination, mutation
- Selection + mutation \Rightarrow continual improvement
- Selection + recombination \Rightarrow innovation
 - fundamental to invention: *generation vs. evaluation*
- Fundamental intuition of GAs: the three work well together

11/2/09 32

Race Between Selection & Innovation: Takeover Time

- Takeover time t^* = average time for most fit to take over population
- Transaction selection: population replaced by s copies of top $1/s$
- s quantifies selective pressure
- Estimate $t^* \approx \ln n / \ln s$

11/2/09 33

Innovation Time

- Innovation time t_i = average time to get a better individual through crossover & mutation
- Let p_i = probability a single crossover produces a better individual
- Number of individuals undergoing crossover = $p_c n$
- Probability of improvement = $p_i p_c n$
- Estimate: $t_i \approx 1 / (p_c p_i n)$


11/2/09 34

Steady State Innovation

- Bad: $t^* < t_i$
 - because once you have takeover, crossover does no good
- Good: $t_i < t^*$
 - because each time a better individual is produced, the t^* clock resets
 - *steady state innovation*
- Innovation number:

$$Iv = \frac{t^*}{t_i} = p_c p_i \frac{n \ln n}{\ln s} > 1$$

11/2/09 35



Other Algorithms Inspired by Genetics and Evolution

- Evolutionary Programming
 - natural representation, no crossover, time-varying continuous mutation
- Evolutionary Strategies
 - similar, but with a kind of recombination
- Genetic Programming
 - like GA, but program trees instead of strings
- Classifier Systems
 - GA + rules + bids/payments
- and many variants & combinations...

11/2/09

37

Additional Bibliography

1. Goldberg, D.E. *The Design of Innovation: Lessons from and for Competent Genetic Algorithms*. Kluwer, 2002.
2. Milner, R. *The Encyclopedia of Evolution*. Facts on File, 1990.



11/2/09

38