

VI
Autonomous Agents
&
Self-Organization

11/4/08 1

Part A
Nest Building

11/4/08 2

Autonomous Agent

- “a unit that interacts with its environment (which probably consists of other agents)
- but acts independently from all other agents in that it does not take commands from some seen or unseen leader,
- nor does an agent have some idea of a global plan that it should be following.”
—Flake (p. 261)

11/4/08 3

Nest Building by Termites
(Natural and Artificial)

11/4/08 4

Resnick’s Termites
 (“Turmites”)

11/4/08 5

Basic procedure

- Wander randomly
- If you are not carrying anything and you bump into a wood chip, pick it up.
- If you are carrying a wood chip and you bump into another wood chip, put down the woodchip you are carrying

— Resnick, *Turtles, Termites, and Traffic Jams*

11/4/08 6

Microbehavior of Turmites


1. *Search for wood chip:*
 - a) If at chip, pick it up
 - b) otherwise wiggle, and go back to (a)
2. *Find a wood pile:*
 - a) If at chip, it's found
 - b) otherwise wiggle, and go back to (a)
3. *Find an empty spot and put chip down:*
 - a) If at empty spot, put chip down & jump away
 - b) otherwise, turn, take a step, and go to (a)

11/4/08 7

Demonstration

[Run Termites.nlogo](#)

11/4/08 8



Why does the number of piles decrease?

- A pile can grow or shrink
- But once the last chip is taken from a pile, it can never restart
- Is there any way the number of piles can increase?
- Yes, and existing pile can be broken into two

11/4/08 10

More Termites

| Termites | 2000 steps | | 10 000 steps | | |
|----------|------------|-----------|--------------|-----------|----------------|
| | num. piles | avg. size | num. piles | avg. size | chips in piles |
| 1000 | 102 | 15 | 47 | 30 | |
| 4000 | 10 | | 3 | 80 | 240 |

11/4/08 11

Termite-Mediated Condensation

- Number of chips is conserved
- Chips do not move on own; movement is mediated by termites
- Chips preferentially condense into piles
- Increasing termites, increases number of chips in fluid (randomly moving) state
- Like temperature

11/4/08 12

An Experiment to Make the Number Decrease More Quickly

- Problem: piles may grow or shrink
- Idea: protect “investment” in large piles
- Termites will not take chips from piles greater than a certain size
- Result: number decreases more quickly
- Most chips are in piles
- But *never* got less than 82 piles

11/4/08 13

Conclusion

- In the long run, the “dumber” strategy is better
- Although it’s slower, it achieves a better result
- By not protecting large piles, there is a small probability of any pile evaporating
- So the smaller “large piles” can evaporate and contribute to the larger “large piles”
- Even though this strategy makes occasional backward steps, it outperforms the attempt to protect accomplishments


11/4/08 14

Mound Building by *Macrotermes* Termites




11/4/08 15

Structure of Mound



11/4/08 16
figs. from Lüscher (1961)


Construction of Mound



- (1) First chamber made by royal couple
- (2, 3) Intermediate stages of development
- (4) Fully developed nest

11/4/08 17
Fig. from Wilson (1971)

Termite Nests



11/4/08 18

Alternatives to Self-Organization

- Leader
 - directs building activity of group
- Blueprint (image of completion)
 - compact representation of spatial/temporal relationships of parts
- Recipe (program)
 - sequential instructions specify spatial/temporal actions of individual
- Template
 - full-sized guide or mold that specifies final pattern

11/4/08 19

Basic Mechanism of Construction (Stigmergy)

- Worker picks up soil granule
- Mixes saliva to make cement
- Cement contains pheromone
- Other workers attracted by pheromone to bring more granules
- There are also trail and queen pheromones

11/4/08 Fig. from Solé & Goodwin 20

Construction of Royal Chamber

11/4/08 21

Construction of Arch (1)

11/4/08 Fig. from Bonabeau, Dorigo & Theraulaz 22

Construction of Arch (2)

11/4/08 Fig. from Bonabeau, Dorigo & Theraulaz 23

Construction of Arch (3)

11/4/08 Fig. from Bonabeau, Dorigo & Theraulaz 24

Basic Principles

- Continuous (quantitative) stigmergy
- Positive feedback:
 - via pheromone deposition
- Negative feedback:
 - depletion of soil granules & competition between pillars
 - pheromone decay

11/4/08 25

Deneubourg Model

- $H(r, t)$ = concentration of cement pheromone in air at location r & time t
- $P(r, t)$ = amount of deposited cement with still active pheromone at r, t
- $C(r, t)$ = density of laden termites at r, t
- Φ = constant flow of laden termites into system

11/4/08 26

Equation for P (Deposited Cement with Pheromone)

$\partial_t P$ (rate of change of active cement) =
 $k_1 C$ (rate of cement deposition by termites)
 $- k_2 P$ (rate of pheromone loss to air)

$$\partial_t P = k_1 C - k_2 P$$

11/4/08 27

Equation for H (Concentration of Pheromone)

$\partial_t H$ (rate of change of concentration) =
 $k_2 P$ (pheromone from deposited material)
 $- k_4 H$ (pheromone decay)
 $+ D_H \nabla^2 H$ (pheromone diffusion)

$$\partial_t H = k_2 P - k_4 H + D_H \nabla^2 H$$

11/4/08 28


Equation for C (Density of Laden Termites)

$\partial_t C$ (rate of change of concentration) =
 Φ (flux of laden termites)
 $- k_1 C$ (unloading of termites)
 $+ D_C \nabla^2 C$ (random walk)
 $- \gamma \nabla \cdot (CVH)$ (chemotaxis: response to pheromone gradient)

$$\partial_t C = \Phi - k_1 C + D_C \nabla^2 C - \gamma \nabla \cdot (CVH)$$

11/4/08 29

Explanation of Divergence



- velocity field = $\mathbf{V}(x,y) = \mathbf{i}V_x(x,y) + \mathbf{j}V_y(x,y)$
- $C(x,y)$ = density
- outflow rate = $\Delta_x(CV_x) \Delta y + \Delta_y(CV_y) \Delta x$
- outflow rate / unit area = $\frac{\Delta_x(CV_x)}{\Delta x} + \frac{\Delta_y(CV_y)}{\Delta y}$

$$\rightarrow \frac{\partial(CV_x)}{\partial x} + \frac{\partial(CV_y)}{\partial y} = \nabla \cdot CV$$

11/4/08 30


Explanation of Chemotaxis Term

- The termite flow *into* a region is the *negative* divergence of the flux through it
 $-\nabla \cdot \mathbf{J} = -(\partial J_x / \partial x + \partial J_y / \partial y)$
- The flux velocity is proportional to the pheromone gradient
 $\mathbf{J} \propto \nabla H$
- The flux density is proportional to the number of moving termites
 $\mathbf{J} \propto C$
- Hence, $-\gamma \nabla \cdot \mathbf{J} = -\gamma \nabla \cdot (C \nabla H)$

11/4/08

31

Simulation ($T = 0$)




11/4/08

fig. from Solé & Goodwin

32

Simulation ($T = 100$)




11/4/08

fig. from Solé & Goodwin

33

Simulation ($T = 1000$)



11/4/08

fig. from Solé & Goodwin

34

Conditions for Self-Organized Pillars

- Will not produce regularly spaced pillars if:
 - density of termites is too low
 - rate of deposition is too low
- A homogeneous stable state results

$$C_0 = \frac{\Phi}{k_1}, \quad H_0 = \frac{\Phi}{k_4}, \quad P_0 = \frac{\Phi}{k_2}$$

11/4/08

35

NetLogo Simulation of Deneubourg Model

[Run Pillars3D.nlogo](#)

11/4/08

36

Interaction of Three Pheromones

- Queen pheromone governs size and shape of queen chamber (template)
- Cement pheromone governs construction and spacing of pillars & arches (stigmergy)
- Trail pheromone:
 - attracts workers to construction sites (stigmergy)
 - encourages soil pickup (stigmergy)
 - governs sizes of galleries (template)

11/4/08

37




Wasp Nest Building and Discrete Stigmergy

11/4/08

Fig. from Solé & Goodwin

38

Structure of Some Wasp Nests




11/4/08

Fig. from Self-Org. Biol. Sys.

39

Adaptive Function of Nests



11/4/08

Figs. from Self-Org. Biol. Sys.

40

How Do They Do It?



11/4/08

41

Lattice Swarms

(developed by Theraulaz & Bonabeau)

11/4/08

42

Discrete vs. Continuous Stigmergy

- Recall: *stigmergy* is the coordination of activities through the environment
- Continuous* or *quantitative* stigmergy
 - quantitatively different stimuli trigger quantitatively different behaviors
- Discrete* or *qualitative* stigmergy
 - stimuli are classified into distinct classes, which trigger distinct behaviors

11/4/08 43

Discrete Stigmergy in Comb Construction

- Initially all sites are equivalent
- After addition of cell, qualitatively different sites created

11/4/08 Fig. from Self-Org. Biol. Sys. 44

Numbers and Kinds of Building Sites

11/4/08 Fig. from Self-Org. Biol. Sys. 45

Lattice Swarm Model

- Random movement by wasps in a 3D lattice
 - cubic or hexagonal
- Wasps obey a 3D CA-like rule set
- Depending on configuration, wasp deposits one of several types of “bricks”
- Once deposited, it cannot be removed
- May be deterministic or probabilistic
- Start with a single brick

11/4/08 46

Cubic Neighborhood







- Deposited brick depends on states of 26 surrounding cells
- Configuration of surrounding cells may be represented by matrices:

$$\begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \times \begin{bmatrix} 0 & 0 & 0 \\ 1 & \bullet & 0 \\ 0 & 0 & 0 \end{bmatrix} \times \begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

11/4/08 Fig. from Solé & Goodwin 47

Hexagonal Neighborhood

11/4/08 Fig. from Bonabeau, Dorigo & Theraulaz 48



Cubic Examples (3)

Step 50000 *m*
Growth Complexity: 141

Step 205000 *n*
Growth Complexity: 153

11/4/08 Figs. from IASC Dept., ENST de Bretagne. 61

Cubic Examples (4)

11/4/08 Figs. from IASC Dept., ENST de Bretagne. 62

Cubic Examples (5)

11/4/08 Figs. from IASC Dept., ENST de Bretagne. 63

An Interesting Example

- Includes
 - central axis
 - external envelope
 - long-range helical ramp
- Similar to *Apicotermes* termite nest

11/4/08 Fig. from Theraulaz & Bonabeau (1995) 64

Similar Results with Hexagonal Lattice

(a)

(b)

(c)

(d)

(e)


- 20x20x20 lattice
- 10 wasps
- All resemble nests of wasp species
- (d) is (c) with envelope cut away
- (e) has envelope cut away

11/4/08 Fig. from Bonabeau & al., *Swarm Intell.* 65

More Hexagonal Examples

11/4/08 Figs. from IASC Dept., ENST de Bretagne. 66


Effects of Randomness (Coordinated Algorithm)



- Specifically different (i.e., different in details)
- Generically the same (qualitatively identical)
- Sometimes results are fully constrained

11/4/08 Fig. from Bonabeau & al., *Swarm Intell.* 67

Effects of Randomness (Non-coordinated Algorithm)



11/4/08 Fig. from Bonabeau & al., *Swarm Intell.* 68

Non-coordinated Algorithms

- Stimulating configurations are not ordered in time and space
- Many of them overlap
- Architecture grows without any coherence
- May be convergent, but are still unstructured

11/4/08 69

Coordinated Algorithm

- Non-conflicting rules
 - can't prescribe two different actions for the same configuration
- Stimulating configurations for different building stages cannot overlap
- At each stage, “handshakes” and “interlocks” are required to prevent conflicts in parallel assembly


11/4/08 70

More Formally...







- Let $C = \{c_1, c_2, \dots, c_n\}$ be the set of local stimulating configurations
- Let (S_1, S_2, \dots, S_m) be a sequence of assembly stages
- These stages partition C into mutually disjoint subsets $C(S_p)$
- Completion of S_p signaled by appearance of a configuration in $C(S_{p+1})$

11/4/08 71

Example



11/4/08 Fig. from Camazine & al., *Self-Org. Biol. Sys.* 72



Conclusions

- Simple rules that exploit discrete (qualitative) stigmergy can be used by autonomous agents to assemble complex, 3D structures
- The rules must be non-conflicting and coordinated according to stage of assembly
- The rules corresponding to interesting structures occupy a comparatively small region in rule-space

11/4/08

Part 6B

79