

Animation of Neuron

- An animated film about nicotine addiction
- A good visualization of a single neuron
- ©2006, Hurd Studios
- Winner of NSF/AAAS Visualization Challenge


View Flash Video


3/30/17


Grey Matter vs. White Matter


(fig. from Carter 1998)


9


























































Intracortical Connections • Dendrites extend 2–4 minicol. diameters • Axons extend 5 × (or even 30–40 × minicol. diameter • Periodic spacing of axon terminal clusters causes entrainment • ~ 2 × 10⁷ connections to macrocolumn 330/17
























Input S	Signals
 Excitatory about 85% of inputs AMPA channels, opened by glutamate Inhibitory about 15% of inputs GABA channels, opened by GABA produced by inhibitory interneurons 	Leakage potassium channels Synaptic efficacy: net effect of: presynaptic neuron to produce neurotransmitter postsynaptic channels to bind it


Membrane Potential (Variables)


- g_e = excitatory conductance
- E_e = excitatory potential ($\sim 0 \text{ mV}$)
- $g_i = \text{inhibitory conductance}$
- E_i = inhibitory potential (–70 mV)
- g_l = leakage conductance
- E_l = leakage potential
- V_m = membrane potential
- θ = threshold


3/30/17




















Rate Code Approximation

- g_e^{θ} is the conductance when $V_m = \theta$
- Rate is a nonlinear function of relative conductance
- What is f?

 $\theta = \frac{g_e^{\theta} E_e + g_i E_i + g_l E_l}{g_e^{\theta} + g_i + g_l}$

 $g_e^{\theta} = \frac{g_i(E_i - \theta) + g_l(E_l - \theta)}{\theta - E_e}$

 $y = f(g_e - g_e^{\theta})$

3/30/17

Activation Function

- · Desired properties:
 - threshold (~0 below threshold)
 - saturation
 - smooth
- Smooth by convolution with Gaussian to account for noise
- Activity update:

$$y_{t+1} = y_t + C(y - y_t)$$

3/30/17



